

Long-term no-till and stover retention each decrease the global warming potential of irrigated continuous corn

VIRGINIA L. JIN¹ , MARTY R. SCHMER¹, CATHERINE E. STEWART², AARON J. SINDELAR^{1,3}, GARY E. VARVEL^{1*} and BRIAN J. WIENHOLD¹

¹Agroecosystem Management Research, United States Department of Agriculture-Agricultural Research Service (USDA-ARS), University of Nebraska-Lincoln, 251 Filley Hall, Lincoln, NE 68583-0937, USA, ²Soil Management and Sugarbeet Research, USDA-ARS, Suite 320, 2150 Centre Avenue, Building D, Fort Collins, CO 80526-8119, USA, ³Monsanto Company, 10975 NW 121st Pl., Malcolm, NE 68402, USA

Abstract

Over the last 50 years, the most increase in cultivated land area globally has been due to a doubling of irrigated land. Long-term agronomic management impacts on soil organic carbon (SOC) stocks, soil greenhouse gas (GHG) emissions, and global warming potential (GWP) in irrigated systems, however, remain relatively unknown. Here, residue and tillage management effects were quantified by measuring soil nitrous oxide (N₂O) and methane (CH₄) fluxes and SOC changes (Δ SOC) at a long-term, irrigated continuous corn (*Zea mays* L.) system in eastern Nebraska, United States. Management treatments began in 2002, and measured treatments included no or high stover removal (0 or 6.8 Mg DM ha⁻¹ yr⁻¹, respectively) under no-till (NT) or conventional disk tillage (CT) with full irrigation ($n = 4$). Soil N₂O and CH₄ fluxes were measured for five crop-years (2011–2015), and Δ SOC was determined on an equivalent mass basis to ~30 cm soil depth. Both area- and yield-scaled soil N₂O emissions were greater with stover retention compared to removal and for CT compared to NT, with no interaction between stover and tillage practices. Methane comprised <1% of total emissions, with NT being CH₄ neutral and CT a CH₄ source. Surface SOC decreased with stover removal and with CT after 14 years of management. When Δ SOC, soil GHG emissions, and agronomic energy usage were used to calculate system GWP, all management systems were net GHG sources. Conservation practices (NT, stover retention) each decreased system GWP compared to conventional practices (CT, stover removal), but pairing conservation practices conferred no additional mitigation benefit. Although cropping system, management equipment/timing/history, soil type, location, weather, and the depth to which Δ SOC is measured affect the GWP outcomes of irrigated systems at large, this long-term irrigated study provides valuable empirical evidence of how management decisions can impact soil GHG emissions and surface SOC stocks.

Keywords: conventional tillage, global warming potential, greenhouse gas intensity, methane, nitrous oxide, no-till, soil organic carbon, stover removal

Received 31 August 2016; revised version received 4 December 2016 and accepted 6 December 2016

Introduction

The majority of global agricultural land base expansion during the last 50 years has been due to increases in irrigated agriculture, which has contributed >40% of increased food production worldwide (FAO, 2011). In the United States, intensification of agricultural systems through irrigation has provided about half the total value of all national crop sales on just 17% of the total U.S. agricultural land base (NASS, 2014; ERS, 2015). These high-yielding production systems, particularly irrigated corn (*Zea mays* L.), are targeted for crop residue removal to supply feedstock for second-generation bioenergy or livestock uses. Retention of corn residues

(e.g., stover) is a recommended conservation practice to reduce soil erosion risk and support fertility by returning organic matter and nutrients to the soil (Wilhelm *et al.*, 2007; Johnson *et al.*, 2014; Karlen *et al.*, 2014). High levels of stover retained on the soil surface, however, can decrease grain yield by interfering with planting, decreasing germination, and increasing disease incidence (Verma *et al.*, 2005; Sindelar *et al.*, 2013). Partial removal of stover from less erosion-prone landscape areas, therefore, could be used to meet livestock or bioenergy feedstock demands while also maintaining or enhancing grain yield in warm temperate environments (Sindelar *et al.*, 2013; Karlen *et al.*, 2014; Wortmann *et al.*, 2016). Pairing stover removal practices with no-till (NT) is expected to further reduce potential risks of soil erosion and soil organic carbon (SOC) loss (Wilhelm *et al.*, 2007; Blanco-Canqui & Lal, 2009; Johnson *et al.*, 2014).

Correspondence: Virginia L. Jin, tel. +1 402 472 5137, fax +1 402 472 0516, e-mail: virginia.jin@ars.usda.gov

*Retired.

The magnitude and direction of SOC change and soil greenhouse gas (GHG) fluxes under irrigated management are variable, leading to uncertainty in global warming potential (GWP) predictions of stover and tillage practices. Higher grain yields under more intensive management can result in greater input-use efficiencies, thereby maintaining or even decreasing GWP relative to less intensively managed systems (Verma *et al.*, 2005; Snyder *et al.*, 2009; Venterea *et al.*, 2011; Grassini & Cassman, 2012). Further, irrigation-induced increases in grain yield and crop biomass relative to non- or deficit irrigation (Norwood, 1999; Maharjan *et al.*, 2014) have been proposed to increase SOC under irrigation, particularly in NT systems (Lal *et al.*, 1998). Recent studies, however, show that irrigated corn production systems have low potential to sequester C because high biomass production is offset by SOC losses from decomposition stimulated by irrigation, especially when the entire soil profile is considered (Verma *et al.*, 2005; Follett *et al.*, 2013; Schmer *et al.*, 2014a).

Although the potential for SOC storage is emphasized frequently as the major factor determining whether a production system will be a net GHG source or sink, SOC changes resulting from residue and tillage management practices could be offset or exacerbated by increased emissions of non-carbon dioxide (CO₂) GHGs such as nitrous oxide (N₂O) and/or methane (CH₄) (Li *et al.*, 2005; Grandy *et al.*, 2006; Mutegi *et al.*, 2010; Powlson *et al.*, 2011; Chen *et al.*, 2013; Schmer *et al.*, 2014a). Some researchers suggest that soil N₂O emissions are expected to be greater in NT compared to conventional tillage (CT) due to greater expected soil water-filled pore space (WFPS) and less aeration due to greater soil bulk densities (Linn & Doran, 1984; Baggs *et al.*, 2003; Rochette, 2008), but others predict smaller emissions under NT because of improved soil structure and cooler soil temperatures (Snyder *et al.*, 2009; Venterea *et al.*, 2011; Sainju *et al.*, 2012; Guzman *et al.*, 2015). Similarly, crop residue management can result in contrasting responses, with residue removal decreasing (Jin *et al.*, 2014; Guzman *et al.*, 2015) or increasing (Congreaves *et al.*, 2016; Lehman & Osborne, 2016) direct GHG emissions from soils.

Because soil changes in response to management can take several years to detect, the ability to quantify management changes improves with experiment duration (Follett *et al.*, 2013; Lehtinen *et al.*, 2014; Jin *et al.*, 2015). Studies that empirically measure soil GHG emissions, SOC stock changes, and energy inputs over a longer time frame (>3 years) are uncommon, and system GWP calculated for many studies rely on estimates or modeled values which are rarely verified for specific management systems in specific geographical locations.

Here, we quantify the *in situ* impacts of continuous crop residue removal and tillage on soil GHG emissions, SOC stocks, and resulting GWP in a long-term irrigated continuous corn system in eastern Nebraska.

Materials and methods

Site description

The study was located at the University of Nebraska–Lincoln Agricultural Research and Development Center, Ithaca, NE (41°9'43.3"N, 96°24'41.4"W; 349 m asl). Soils are silt loams of the Tomek (fine, smectitic, mesic Pachic Argiudoll) and Filbert (fine, smectitic, mesic Vertic Argialboll) series, 0–2% slopes (Soil Survey Staff, 2014). The 30-year mean air temperature and precipitation are 9.8 °C and 74 cm, respectively (1981–2010) (NCDC; Station ID Mead 6 S). Historically, this area was cropped with corn (*Z. mays* L.), soybean [*Glycine max* (L.) Merr.], oat (*Avena sativa* L.), and alfalfa (*Medicago sativa* L.).

Continuous corn was established at the site in 2000. Details of the study design are reported in Schmer *et al.* (2014a). Briefly, the experiment is a randomized complete block design ($n = 6$ blocks) with factorial treatments in split plots. Tillage is the whole-plot factor (NT, CT; 9 m × 45.6 m) and stover removal rate is the subplot factor (none, 0%; medium, ~35%; high, ≥70%; 9 m × 15.2 m). All treatment combinations were in place by 2002. Nitrogen fertilizer was side-dressed at V4–V6 stages at varying rates from 2001 to 2006. Since 2007, N fertilizer has been side-dressed as granular urea (46-0-0) at 202 kg N ha⁻¹ yr⁻¹ using a 12-row applicator with injector knives placing N between rows to a soil depth of 10–15 cm. The disk treatment area is tilled each year, usually just before planting in spring, to a depth of 15–20 cm. Irrigation was applied with a solid-set sprinkler until 2002, then with a linear-move irrigation system since that time. Annual irrigation amounts ranged from 25 to 266 mm from 2001 to 2015, with a 14-year average of 116 ± 17 mm. Corn stover is removed in the fall after grain harvest using a flail chopper set to a 10-cm cutting height.

Crop productivity

Plant production values, soil GHG emissions, and soil organic C changes are reported here for only the 'none' and 'high' stover removal levels under NT and CT in four of the six blocks where GHG emissions were measured. Annual aboveground dry matter (DM) production was measured by hand from a 0.76 m × 3.04 m area shortly after physiological maturity (September or October) each year. Ears were removed, and then stalks were cut at ground level, chopped, and weighed, and a subsample was dried at 60 °C until constant mass. Ears were dried, weighed, and shelled to calculate DM grain yields. Cob weights were added to the DM stalk biomass to calculate total nongrain DM (i.e., stover) production. Harvest index (HI) was calculated as the ratio of grain to total aboveground biomass production.

Soil greenhouse gas emissions

Soil GHG emissions were sampled from four of six treatment blocks from April 2011 through May 2016 for a total of 100 sampling events that spanned five full crop-years (i.e., >1500 individual flux observations). In April 2011, four blocks were selected at random for intensive soil GHG measurement using static vented chambers. The same four blocks have been sampled over time as per the USDA-ARS standardized protocol for the Greenhouse gas Reduction through Agricultural Carbon Enhancement network (GRACEnet) (Hutchinson & Mosier, 1981; Parkin & Venterea, 2010). Vented gas sampling chambers were covered with reflective insulation, and both chambers and sampling bases that were installed in the field were constructed of 20-gage stainless steel. Each base covered an area of 1707 cm² (52.7 cm × 32.4 cm) and was installed to a soil depth of 5–7 cm, resulting in a base height of 5–7 cm above ground. One chamber base was placed well within each treatment plot and oriented perpendicularly to rows such that the short edge was parallel to the row and the remainder of the base extended into the between-row area. The base footprint approximated a 1 : 2 ratio of within-row to between-row soil microsites. The between-row microsite included the injection furrow where urea fertilizer was applied. Bases were removed and re-installed throughout the growing season to accommodate field management operations (e.g., tillage, planting, fertilization, harvest). After any reinstallation, bases were allowed to equilibrate for at least 24 h (but usually 2–4 days) before gas sampling to minimize immediate soil disturbance effects (Reicosky *et al.*, 2005).

At each sampling event, headspace gas samples were collected with syringes then injected into evacuated vials at four evenly spaced time points over 30 min (0, 10, 20, and 30 min). To account for diurnal variability and also approximate daily average temperature, sampling was conducted early to mid-morning on each sampling date. Sampling occurred every 5–10 days during the growing season (May to September), with greater frequency following tillage operations and N fertilizer application. Nongrowing season measurements were collected monthly as weather and ground conditions allowed.

Ancillary measurements at each sampling date included air temperature, soil temperature at 15 cm, and soil moisture from 0 to 15 cm depth as measured by a handheld time domain reflectometer (FieldScout TDR 300; Spectrum Technologies, Aurora, IL, USA) with a site-specific calibration. Volumetric soil water content was converted to water-filled pore space (WFPS) using soil bulk density values.

Sample GHG concentrations were measured within 10 d of collection using a headspace autosampler (CombiPAL; CTC Analytics, Zwingen, Switzerland) connected to a gas chromatograph (450-GC; Varian, Middelburg, the Netherlands) equipped with separate detectors for the simultaneous measurement of N₂O (electron capture detector), CH₄ (flame ionization detector), and CO₂ (thermal conductivity detector). Soil CO₂ data are not presented here. Soil N₂O and CH₄ fluxes were calculated as a linear or quadratic change in headspace gas concentration over time within the enclosed chamber volume (Wagner *et al.*, 1997; Venterea *et al.*, 2011), and soil N₂O fluxes were corrected for suppression of the

surface–atmosphere concentration gradient (Venterea, 2010). Fluxes were considered nonzero if rates were greater than (i.e., production) or less than (i.e., consumption) the flux detection limits calculated on the basis of analytical precision, chamber deployment time, and ambient gas concentration (Parkin *et al.*, 2012). Area-scaled N₂O and CH₄ emissions were estimated by linear interpolation of flux rates between sampling dates, then summing daily rates over each growing season (i.e., trapezoidal integration method). Yield-scaled N₂O emissions, or greenhouse gas intensity (GHGI), were calculated by dividing area-scaled N₂O emissions by DM grain yield. Total non-CO₂ GHG emissions were calculated by summing area-based annual fluxes of N₂O and CH₄ after converting to CO₂ equivalents (Mg CO₂ eq ha⁻¹) using GWP values of 298 and 25, respectively (Forster *et al.*, 2007).

Soil organic carbon change and system global warming potential

Soil samples for SOC and GWP calculations were collected in May 2001, November 2010, and November 2014. In 2001 and 2010, four cores (3.5 cm diameter) were sampled from each subplot and composited by depth increment (0–15, 15–30, 30–60, 60–90, 90–120, 120–150 cm) (Schmer *et al.*, 2014a). In 2014, two cores (4.1 cm diameter) were sampled from each subplot and composited by depth increment (0–7.5, 7.5–15, 15–30 cm). Soil bulk densities were determined for each soil depth increment using core volumes and dry weights (dried at 105 °C) for soils collected in all years. A subsample of fresh soil from each increment was passed through a 2-mm sieve, oven-dried at 55 °C (up to 48 h), ground to pass a 150-mm screen, and analyzed for total C and N concentrations (Follett & Pruessner, 2001). Surface soils were carbonate-free.

Due to differences in the time of year and depth increments sampled between years, changes in SOC stocks were calculated on an equivalent mass basis that approximated a soil depth of 30 cm (Lee *et al.*, 2009). Changes in SOC (Δ SOC; Mg CO₂ eq ha⁻¹) were calculated for the same four experimental blocks measured for non-CO₂ GHG fluxes. Changes were calculated for three time periods: (i) 2001–2010 (early period), (ii) 2010–2014 (later period), and (iii) 2001–2014 (full period of study).

Global warming potential was calculated for each management system as total non-CO₂ soil GHG emissions plus total fuel usage from agronomic operations (Table S1) minus total SOC stock changes for each early and later time period (Mosier *et al.*, 2006; Cavigelli & Parkin, 2012). Total non-CO₂ GHG emissions for 2001–2010 were estimated for each management scenario (NT-None, NT-High, CT-None, CT-High) by multiplying the number of crop-years ($n = 10$) by the mean annual emissions measured from 2011 to 2015. Total non-CO₂ GHG emissions for 2010–2014 were calculated as the total emissions measured over those crop-years. Total non-CO₂ GHG emissions for the full study period were calculated as the sum of the early and later periods. Fuel usage was estimated from other studies (Adler *et al.*, 2007; Schmer *et al.*, 2014b), converted to CO₂ equivalents per hectare using the most recent carbon intensity values for diesel (102.82 kg CO₂ eq GJ⁻¹) or electricity (105.62 kg CO₂ eq GJ⁻¹) (California

Air Resources Board, 2014), and assumed to be constant across all years (Table S1). A positive GWP value indicated that the system was a GHG source, and negative GWP value indicated that the system was a GHG sink. To standardize units for time, Δ SOC and GWP are presented on an annual basis ($\text{Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$).

Statistical analyses

Soil temperature, soil WFPS, aboveground biomass production, grain yield, HI, stover removal, area-scaled GHG emissions (N_2O , CH_4 , total non- CO_2 GHGs), and GHGI was analyzed using the GLIMMIX procedure of SAS (SAS Institute, 2014). Tillage (T), stover removal (R), and their interactions were considered fixed effects, with year (Y) and block as random effects. For analyses including year, the covariance structure for each response variable was selected based on the smallest Akaike information criterion (Littell *et al.*, 2006; Loughin, 2006). Five-year total emissions of non- CO_2 GHGs (2011–2016) and Δ SOC and GWP for each time period (2001–2010, 2010–2014, 2001–2014) were analyzed using the GLIMMIX procedure with T, R, and $T \times R$ considered as fixed effects and block as a random effect. All treatment effects were considered significant at $P \leq 0.05$. Data were ln-transformed for normality when necessary. Multiple comparisons between significant treatment responses were evaluated with Bonferroni-adjusted P -values using the SLICE option in the LSMEANS statement ($P \leq 0.05$). Values are reported as means \pm standard error (SE). All plot-level data are available online (USDA-ARS Data Portal, <http://nrrc.ars.usda.gov/arsdataportal/>) (Del Grosso *et al.*, 2013).

Results

Environmental conditions

Average air temperature and rainfall conditions varied widely across the 2011–2015 crop-years (Fig. 1a, b). Mean annual air temperature ranged from 9.1 to 11.4 °C, and the 5-year mean (10.5 °C) was $7 \pm 5\%$ above the preceding 30-year average (9.8 °C). Mean annual precipitation inputs ranged from 43.2 to 82.2 cm, and the 5-year mean (63 cm) was $14 \pm 10\%$ below the preceding 30-year average (74 cm) primarily due to two drought years (2012, 2013). Growing season air temperatures ranged from 20.7 to 23.1 °C, and irrigation inputs ranged from 2.5 to 16.6 cm per year.

Measured soil temperatures (15 cm depth) were greater in CT (15.7 ± 0.3 °C) compared to NT (15.3 ± 0.3 °C) across all 100 sampling dates during 2011–2015 crop-years ($P_T = 0.0334$). Measured soil WFPS (0–15 cm depth) was less in CT ($45.7 \pm 0.3\%$) compared to NT ($50.9 \pm 0.3\%$) across all sampling dates, and the difference between CT and NT was greatest during the 2012 drought year (40.5% and

47.9%, respectively) ($P_{Y \times T} = 0.0010$). There were no main or interaction effects of stover removal on either soil temperature or soil WFPS.

Crop responses

Total aboveground production (grain + nongrain biomass, $\text{Mg DM ha}^{-1} \text{ yr}^{-1}$) and grain yield differed across the 2011–2015 crop-years ($P_Y < 0.005$), but were not affected by tillage or stover removal treatments (Tables 1 and 2). Total aboveground production was least in 2015, greatest in 2013, and averaged $18.17 \pm 0.74 \text{ Mg DM ha}^{-1} \text{ yr}^{-1}$ across the five crop-years (Table 2). Grain yields were numerically smallest in 2012, greatest in 2013, and averaged $9.85 \pm 0.46 \text{ Mg DM ha}^{-1} \text{ yr}^{-1}$ across the five crop-years (Table 2). Harvest index (HI) was marginally greater when stover was removed ($\text{HI}_{\text{High}} = 0.547$) compared to no removal ($\text{HI}_{\text{None}} = 0.536$) ($P_R = 0.0817$) (Table 1). Harvest index values were $\text{NT} > \text{CT}$ in 2011 and 2015, but $\text{NT} < \text{CT}$ in the 2012 drought year ($P_{Y \times T} < 0.005$) (Table 2). Values for HI ranged from 0.47 to 0.60 under NT and from 0.48 to 0.61 under CT. The quantity of stover biomass removed was not affected by tillage but differed by year ($P_Y < 0.0001$; Table 1), ranging from 4.90 in 2014 to 8.84 Mg DM ha^{-1} in 2011. The mean stover biomass removed over the five crop-years was $6.87 \pm 0.64 \text{ Mg DM ha}^{-1}$ and was equivalent to 84% of nongrain aboveground biomass.

Soil N_2O fluxes and greenhouse gas intensity

Measured daily soil N_2O fluxes ranged from -1.3 to $519.6 \text{ g N}_2\text{O-N ha}^{-1} \text{ day}^{-1}$, with a median value of $2.2 \text{ g N}_2\text{O-N ha}^{-1} \text{ day}^{-1}$ and mean value of $14.6 \pm 1.0 \text{ g N}_2\text{O-N ha}^{-1} \text{ day}^{-1}$ over the 2011–2015 crop-years ($n = 1573$) (Figs 1c and 2). Eighty-four percent of all observations were nonzero fluxes. The greatest daily N_2O fluxes occurred one to 3 weeks following fertilizer application (Fig. 1). Peak fluxes occurred when soil temperature > 25 °C and soil WFPS $> 60\%$ (Fig. 2). Smaller N_2O efflux events occurred in February to March, corresponding with spring thawing of frozen soils. Although soils generally emitted N_2O , 2% of all nonzero observations were negative fluxes, indicating soil N_2O consumption.

Area-scaled annual N_2O emissions ranged from 0.4 to 7.2 $\text{kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$ over all treatments, with a mean annual value of $3.2 \pm 0.2 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$ for the 2011–2015 crop-years (Fig. 3a, d). Annual N_2O emissions did not differ between years in NT ($2.9 \pm 0.1 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$), but emissions were 42% and 73% greater in CT than in NT for the 2013 and 2014 crop-years, respectively (Fig. 3a) ($P_{Y \times T} = 0.0174$).




Fig. 1 (a) Daily air temperature ($^{\circ}\text{C}$) with growing season mean daily air temperature (GS_T); (b) water inputs (mm day^{-1}) from precipitation (gray bars) and irrigation (black bars) with annual precipitation total, growing season precipitation total ($\text{GS}_{\text{Precip}}$), and total irrigation inputs; and effects of tillage (NT, CT) and stover removal (none, high) on soil fluxes of (c) N_2O ($\text{g N ha}^{-1} \text{ day}^{-1}$) and (d) CH_4 ($\text{g C ha}^{-1} \text{ day}^{-1}$) for 2011–2015 crop-years. Arrows in (c) indicate N fertilizer application dates for each year.

Annual N_2O emissions were less when stover was removed ($2.8 \pm 0.2 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$) than the no stover removed treatment (i.e., stover retained) ($3.5 \pm 0.2 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$) ($P_R = 0.0311$) (Fig. 3d).

Yield-scaled annual N_2O emissions (GHGI) ranged from 0.034 to $1.081 \text{ kg N}_2\text{O-N Mg}^{-1}$ grain over all treatments, with a mean annual value of $0.333 \pm 0.019 \text{ kg N}_2\text{O-N Mg}^{-1}$ DM grain for the 2011–2015 crop-years (Fig. 4a, b). Over all crop-years, GHGI with NT ($0.299 \pm 0.021 \text{ kg N}_2\text{O-N Mg}^{-1}$ DM grain) was less than CT ($0.363 \pm 0.027 \text{ kg N}_2\text{O-N Mg}^{-1}$ DM grain) ($P_T = 0.0368$), and GHGI was less when stover

was removed ($0.289 \pm 0.022 \text{ kg N}_2\text{O-N Mg}^{-1}$ DM grain) compared to retained ($0.373 \pm 0.020 \text{ kg N}_2\text{O-N Mg}^{-1}$ DM grain) ($P_R = 0.0107$).

Soil CH_4 fluxes

Measured daily soil CH_4 fluxes ranged from -23.0 to $97.0 \text{ g CH}_4\text{-C ha}^{-1} \text{ day}^{-1}$, with a median value of $-0.3 \text{ g CH}_4\text{-C ha}^{-1} \text{ day}^{-1}$ and a mean value of $0.2 \pm 0.1 \text{ g CH}_4\text{-C ha}^{-1} \text{ day}^{-1}$ over the 2011–2015 crop-years ($n = 1539$) (Fig. 1d). Eighty-four percent of all observations were nonzero fluxes. Of all nonzero

Table 1 Significance of *F* values for fixed sources of variation for total aboveground production (nongrain + grain, grain yield, harvest index, and stover removed for 2011–2015 crop-years

Source of variation	df	Total aboveground production	Grain yield	Harvest index	Stover removed
		Pr > <i>F</i>			
Year (Y)	4	0.0040*	0.0017	<0.0001	<0.0001
Tillage (T)	1	0.6914	0.6383	0.4977	0.5833
Removal (R)	1	0.9228	0.4278	0.0817	<0.0001
Y × T	4	0.2890	0.1422	0.0049	0.3139
Y × R	4	0.2898	0.4271	0.2349	<0.0001
T × R	1	0.0798	0.0977	0.3505	0.5833
Y × T × R	4	0.3900	0.2805	0.2192	0.3139

*Bold text indicates significant *P*-values ($P \leq 0.05$).

Table 2 Total aboveground production, grain yield, and harvest index in response to different soil management and stover removal practices for 2011–2015 crop-years

	No-till			Conventional tillage			
Removal level	None	High	Mean	None	High	Mean	Annual mean
Total aboveground production, Mg DM ha ⁻¹							
2011	18.37	19.70	19.04	18.55	19.65	19.10	19.07b†
2012	17.24	18.43	17.84	20.99	14.71	17.85	17.84ab
2013	20.33	18.87	19.60	21.40	20.69	21.05	20.32b
2014	16.57	18.63	17.60	17.26	18.29	17.77	17.69ab
2015	16.56	18.58	17.57	14.15	14.33	14.24	15.91a
Mean	17.81	18.84	18.33	18.47	17.53	18.00	18.17
Grain yield, Mg DM ha ⁻¹							
2011	9.54	10.12	9.83	8.67	9.45	9.06	9.44b
2012	7.88	9.16	8.52	10.89	7.69	9.29	8.91ab
2013	11.18	10.57	10.88	11.45	11.70	11.58	11.23b
2014	9.38	11.72	10.54	10.49	11.05	10.77	10.65ab
2015	9.81	10.41	10.11	7.97	7.92	7.94	9.03a
Mean	9.55	10.40	9.98	9.90	9.56	9.73	9.85
Harvest index‡							
2011	0.52	0.51	0.52AB*	0.47	0.48	0.48A	0.50
2012	0.45	0.50	0.47A*	0.52	0.52	0.52AB	0.50
2013	0.55	0.56	0.56BC	0.54	0.57	0.55BC	0.55
2014	0.56	0.63	0.60C	0.61	0.61	0.61D	0.60
2015	0.59	0.56	0.58C*	0.55	0.54	0.54C	0.56
Mean	0.54	0.55	0.54	0.54	0.54	0.54	0.54

*Tillage difference within a given year ($P \leq 0.05$).

†Different lowercase letters and bold text indicate annual differences in aboveground production and yield ($P \leq 0.05$).

‡Different uppercase letters and bold text indicate annual differences within tillage treatments for HI.

CH₄ fluxes, 44% of observations indicated CH₄ consumption (i.e., negative flux). There were no clear seasonal patterns of CH₄ fluxes over the five crop-years.

Area-scaled annual CH₄ fluxes ranged from -0.67 to 1.28 kg CH₄-C ha⁻¹ yr⁻¹ over all treatments, averaging 0.19 ± 0.04 kg CH₄-C ha⁻¹ yr⁻¹ for the 2011–2015

crop-years (Fig. 3b, e). Annual CH₄ fluxes did not differ between years in NT (0.07 ± 0.06 kg CH₄-C ha⁻¹ yr⁻¹) and were not different from zero flux in four of five crop-years. In contrast, CT soils emitted CH₄ in three of five crop-years, and annual CH₄ emissions were 399%, 89%, and 261% greater than NT soils in 2012, 2013, and 2015, respectively ($P_{Y \times T} = 0.0024$).




Fig. 2 Effects of soil temperature (soil T, °C) and soil water-filled pore space (WFPS, %) on daily N_2O fluxes ($\text{g N ha}^{-1} \text{ day}^{-1}$) for 2011–2015 crop-years.

(Fig. 3b). There was no stover removal effect in CT ($0.31 \pm 0.13 \text{ kg CH}_4\text{-C ha}^{-1} \text{ yr}^{-1}$), but NT soils emitted CH_4 in the high stover removal treatment ($0.17 \pm 0.06 \text{ kg CH}_4\text{-C ha}^{-1} \text{ yr}^{-1}$) and were CH_4 neutral when no stover was removed ($-0.03 \pm 0.08 \text{ kg CH}_4\text{-C ha}^{-1} \text{ yr}^{-1}$) ($P_{\text{T} \times \text{R}} = 0.0161$) (Fig. 3e).

Total emissions of non- CO_2 gases

Area-scaled, total annual emissions of non- CO_2 GHGs (N_2O plus CH_4 ; $\text{Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) was dominated by N_2O (>99%) (Fig. 3c, f). The mean annual GHG emission for all treatments over the five crop-years was $3.0 \pm 0.2 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$. The majority of emissions occurred during the growing season (81–93%). Total emissions did not differ between years in NT ($2.7 \pm 0.1 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$), but CT > NT for 2013 and 2014 crop-years ($P_{\text{Y} \times \text{T}} = 0.0100$). Total emissions under high stover removal ($2.6 \pm 0.2 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) were less than no removal ($3.3 \pm 0.2 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) ($P_{\text{R}} = 0.0325$). There were no other treatment interaction effects for year, tillage, and stover removal on total annual non- CO_2 GHG emissions. The 5-year cumulative emissions did not differ between CT or NT, but was 23% less in the high stover removal treatment ($12.2 \pm 1.5 \text{ Mg CO}_2 \text{ eq ha}^{-1}$) compared to no removal ($15.8 \pm 1.4 \text{ Mg CO}_2 \text{ eq ha}^{-1}$) ($P_{\text{R}} = 0.0513$). Values for the mean annual total emissions ($\text{Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) used to estimate GWP for the early study period (2001–2010) were 2.8, 2.1, 3.5, and 2.8 for NT-None, NT-High, CT-None, and CT-High treatments, respectively.

Soil organic carbon changes

Soil organic carbon stocks (ΔSOC) in surface soils (fixed mass to ~30 cm soil depth) decreased over time in all management systems. Relative to 2001 SOC values, 2014 SOC changes were not different than zero for NT-None, marginal SOC loss under CT-None (−5%; $P = 0.1327$), and significant SOC losses under both NT-High (−9%) and CT-High (−10%) ($P \leq 0.01$). For all management periods, however, there were no interaction effects of stover and tillage management (Figs 5 and S1). Conservation practices of stover retention and NT resulted in no SOC change, but 9% of SOC was lost with a high level of stover removal and 7% of SOC was lost with CT over the 14-year study period (Fig. S1).

During the early study period (2001–2010), annual ΔSOC was affected only by stover management ($P_{\text{R}} = 0.0100$), with SOC loss occurring with stover removal ($-1.4 \pm 0.2 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) (Fig. 5c). Although there were no tillage differences for annual ΔSOC , SOC loss rates were less than zero for both NT ($-0.6 \pm 0.4 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) and CT ($-1.0 \pm 0.3 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) (Fig. 5a). During the later study period (2010–2014), there were no main or interaction effects of tillage or stover management on annual ΔSOC , and although values were not different than zero, all losses were numerically greater than annual ΔSOC measured in the early period (2001–2010). Over the full study period (2001–2014), annual ΔSOC was affected by stover removal ($P_{\text{R}} = 0.0062$) with annual SOC losses occurring only when stover was removed ($-1.6 \pm 0.3 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$). Although there were no tillage differences over the full study period, annual ΔSOC was less than zero in CT only ($-1.2 \pm 0.3 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$), indicating SOC loss.

Global warming potential

Annual GWP values were greater than zero for all management systems for all time periods, indicating that all systems were net GHG sources (Fig. 5b, d). During the early study period (2001–2010), annual GWP was affected by both tillage ($P_{\text{T}} = 0.0157$) and stover removal ($P_{\text{R}} = 0.0299$). Global warming potential was greater in CT ($4.9 \pm 0.3 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) compared to NT ($3.8 \pm 0.4 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$), and when stover was removed ($4.9 \pm 0.2 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) compared to retained ($3.9 \pm 0.3 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$). During the later period of the study (2010–2014), annual GWP was not affected by either tillage or stover removal, but were numerically greater in all systems compared to the earlier study period (mean GWP for all systems $5.2 \pm 0.9 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$).




Fig. 3 Effects of tillage (a–c) and stover removal (d–f) on soil fluxes of N₂O (kg N ha^{−1} yr^{−1}), CH₄ (kg C ha^{−1} yr^{−1}), and total non-CO₂ GHGs (Mg CO₂ eq ha^{−1} yr^{−1}) for 2011–2015 crop-years. In (a–c), different letters indicate yearly differences; tillage differences by year denoted by **P* ≥ 0.05, ****P* ≥ 0.005. In panels (d–f), asterisks indicate stover differences. In panels (b, e), (†) indicate fluxes different than zero.




Fig. 4 Mean greenhouse gas intensity (kg N₂O-N Mg^{−1} DM grain) for 2011–2015 crop-years by (a) tillage and (b) stover removal rate. Asterisks indicate a treatment difference (**P* ≥ 0.05, ***P* ≥ 0.01).




Fig. 5 Effects of tillage (a, b) and stover removal (c, d) on annual soil organic carbon changes (ΔSOC) and net global warming potential (GWP) ($\text{Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) for 2001–2010, 2010–2014, and the full 2001–2014 time horizon. Negative ΔSOC values indicate SOC losses. Within each time period, treatment differences are denoted by * $P \geq 0.05$, ** $P \geq 0.01$, and $^{\$}P = 0.0817$. In panels (a, c), daggers (†) indicate SOC changes that are different than zero.

Over the full study period (2001–2014), annual GWP was affected by stover removal ($P_R = 0.0309$) and, to a lesser extent, tillage ($P_T = 0.0817$). Global warming potential was greater when stover was removed ($5.1 \pm 0.4 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) compared to retained ($4.1 \pm 0.4 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$), and in CT ($5.2 \pm 0.3 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$) compared to NT ($4.0 \pm 0.4 \text{ Mg CO}_2 \text{ eq ha}^{-1} \text{ yr}^{-1}$).

Discussion

Although crop yields did not differ between management treatments in the last five years of this 14-year irrigated continuous corn study, non- CO_2 GHG emissions, surface SOC changes, and system GWP were affected by the main effects of year, tillage, and/or stover removal. Changes in SOC affected GWP more so than changes in soil non- CO_2 GHG emissions as a result of management. Management differences in SOC changes and soil GHG emissions, however, showed high spatial and temporal variability, emphasizing the importance of long-term measurements in capturing soil responses to management. Our findings suggest that stover management decisions in irrigated systems have more

impact on SOC stocks than tillage management, but that both tillage and stover management decisions affect overall system GWP. Although the absence of management interactions indicated no added benefit in pairing conservation management practices (e.g., NT, stover retention) compared to using each practice individually, other agroecosystem goals such as benefits to soil health and reduced soil erosion risks will also affect producer decisions (Blanco-Canqui & Lal, 2009; Karlen *et al.*, 2011; Stewart *et al.*, 2015).

Crop productivity

Neither tillage nor stover removal affected mean total aboveground biomass nor grain yields over the 2011–2015 crop-years. Stover removal, however, tended to increase HI (Tables 1 and 2). In a previous study at this site, Schmer *et al.* (2014a) found greater grain yields under high stover removal (2001–2010) under NT but not CT. Other irrigated corn studies have also found improved grain yields with stover removal (Sims *et al.*, 1998; Halvorson & Stewart, 2015; Wortmann *et al.*, 2016). Warmer soil temperatures associated with stover removal likely provided more favorable conditions for

corn germination, growth, and grain production (Kaspar *et al.*, 1990; Sims *et al.*, 1998; Blanco-Canqui & Lal, 2007; Sindelar *et al.*, 2013; Halvorson & Stewart, 2015). In contrast, some irrigated corn studies have reported that stover removal did not affect yield (Biau *et al.*, 2013; Huang *et al.*, 2013; Kenney *et al.*, 2013), and progressive yield declines have been found in rainfed corn production sites when stover was removed (Wilhelm *et al.*, 2004; Blanco-Canqui & Lal, 2007).

While the present study found no tillage effects on production since 2010, lower grain yields with NT have been noted in other irrigated corn studies (Sims *et al.*, 1998; Halvorson *et al.*, 2006) as well as in rainfed studies (Grandy *et al.*, 2006; Bundy *et al.*, 2011; Ogle *et al.*, 2012; Sindelar *et al.*, 2015). In these studies, reductions in crop yield with NT were often attributed to cooler early season soil temperatures that potentially slowed early spring development and delayed tasseling. Although we found warmer soil temperatures in CT compared to NT in this study, temperature differences did not result in a CT yield advantage over the last 5 years.

Daily fluxes of soil N₂O and CH₄

For all management systems, daily N₂O emissions rose sharply 1–3 weeks after fertilizer application (Fig. 1). High emissions rates following N fertilizer application is expected, and the length of time to these high fluxes depends on N fertilizer type, application method, and soil conditions (e.g., moisture, temperature) (Halvorson *et al.*, 2008; Venterea *et al.*, 2011; Drury *et al.*, 2012; Decock, 2014; Maharjan *et al.*, 2014). Across all individual N₂O measurements ($n = 1573$), peak daily emissions occurred when soil temperatures were warmer than 25 °C and soil WFPS >60% (Fig. 2). Although there are many biotic and abiotic pathways that potentially contribute to soil N₂O emissions (Butterbach-Bahl *et al.*, 2013), the aerobic process of nitrification is generally considered the primary N₂O source when WFPS is 30–60%, and the anaerobic process of denitrification is the primary N₂O source when WFPS >60% (Linn & Doran, 1984; Davidson, 1991; Bateman & Baggs, 2005). Depending on soil texture, many soils at 60% WFPS are at field capacity where soil oxidative (e.g., nitrification) and reductive (e.g., denitrification) processes co-occur while transitioning in response to soil oxygen levels (Linn & Doran, 1984; Davidson, 1991). In this study, only 7% of all soil observations had WFPS >60%, and these soil moistures were associated with the greatest daily emissions rates measured over the five crop-years, especially when soils were warm. This suggests that episodic peak events during the growing season can heavily influence the total annual emissions of N₂O (Butterbach-Bahl *et al.*, 2013), emphasizing the

importance of an adequate sampling frequency to accurately quantify annual N₂O emissions (Parkin, 2008; Parkin & Venterea, 2010).

The optimum range for N₂O emissions is proposed to occur at 70–80% WFPS (Davidson, 1991; Chen *et al.*, 2013), beyond which N₂O levels are expected to drop as it undergoes complete stepwise denitrification to N₂ (Chapuis-Lardy *et al.*, 2007; Butterbach-Bahl *et al.*, 2013). In a review of European soils, Schaufler *et al.* (2010) found that N₂O emissions declined in very few of even the wettest soils, suggesting that moisture conditions beyond the optimum WFPS is infrequent in upland soils (Butterbach-Bahl *et al.*, 2013). Although we found that 2% of all nonzero N₂O fluxes were negative in this irrigated system study, there was similarly no evidence that N₂O consumption was occurring in the wettest soils. Rather, the majority of N₂O consumption events occurred in early spring (February, March), when median soil temperatures were cool (3.8 °C) and median soil moisture was 52% WFPS. During this time, a minor peak in soil N₂O effluxes also occurred, indicating that both oxidative and reductive processes were active when soils were undergoing freeze–thaw events (Fig. 1). Increases in N₂O effluxes in early spring associated with soil thawing have been measured in both rainfed (Congreaves *et al.*, 2016) and irrigated systems (Ellert & Janzen, 2008; Cui *et al.*, 2012). While the mechanisms for soil N₂O consumption are not yet clearly understood, increased N₂O consumption in cooler soils with more moderate soil moisture content could result from increased nitrifier denitrification, aerobic denitrification, or other shifts in active microbial community or abiotic processes (Chapuis-Lardy *et al.*, 2007; Ellert & Janzen, 2008). Further, in colder climates where soils are frozen for extended periods, nongrowing season emission events can constitute 30–90% of annual N₂O fluxes (van Bochove *et al.*, 2000; Teepe *et al.*, 2000; Congreaves *et al.*, 2016), but was only a minor contribution (~13%) to total annual emissions in this current study.

Upland agricultural soils are typically minor emitters or minor sinks for CH₄ (Bronson & Mosier, 1994; Mosier *et al.*, 2006). Of the 1539 daily CH₄ measurements made in this study, 44% of all nonzero CH₄ fluxes were negative, indicating CH₄ consumption. Although CH₄ consumption is expected to occur in aerobic soils while CH₄ production occurs in anoxic soils (Conrad, 1996), we found no clear seasonal patterns between soil conditions and daily CH₄ fluxes over the five crop-years measured here.

Annual fluxes of non-CO₂ soil greenhouse gases

In this study, we found both area- and yield-scaled soil N₂O emissions were greater when stover was retained

compared to removed and under CT compared to NT, with no interaction between stover and tillage practices. The absence of tillage and residue management interactions on soil GHG emissions has also been noted by others (Baggs *et al.*, 2003; Mahli & Lemke, 2007). Cumulative annual emissions of non-CO₂ soil GHGs differed between years for each tillage treatment such that emissions were greater under CT (Figs 2 and 3).

Area-scaled and yield-scaled (GHGI) annual N₂O emissions reported here were within the range of measured or modeled values for other irrigated corn production systems (Adviento-Borbe *et al.*, 2007; Liu *et al.*, 2011; Cui *et al.*, 2012; Grassini & Cassman, 2012; Aguilera *et al.*, 2013; Huang *et al.*, 2013; Jin *et al.*, 2014; Zhang *et al.*, 2014). Further, both area-scaled and yield-scaled emissions showed similar responses to management treatments, which have also been noted in other irrigated and nonirrigated studies (Drury *et al.*, 2012; Huang *et al.*, 2013; van Kessel *et al.*, 2013). For area-scaled N₂O and CH₄ emissions, mean annual fluxes were relatively stable across the five crop-years under NT, but more variable under CT such that CT soils were stronger GHG emitters in some years (2013, 2014 for N₂O; 2012, 2013, 2015 for CH₄). Notably, NT was CH₄ neutral while CT was a CH₄ source. In addition, stover removal increased CH₄ emissions compared to no removal in NT but not in CT. This contrasts with other irrigated corn studies that showed no tillage effects on soil CH₄ fluxes (Mosier *et al.*, 2006) or greater CH₄ emissions under NT compared to CT (Alluvione *et al.*, 2009). When converted to CO₂ equivalents, however, CH₄ contributed <1% of the total annual non-GHG emissions; thus, the remainder the discussion focuses on N₂O below.

The variable effects of tillage on soil N₂O emissions emphasize how soils, weather, climate, and management duration can affect agroecosystem responses to management. A meta-analysis of 239 direct comparisons between NT/reduced tillage (RT) with CT found no overall tillage differences in soil N₂O emissions, but that in drier climates, NT/RT increased emissions in the short term (<10 years) by 56% but decreased emissions in the longer term (>10 years) by 27% compared to CT (van Kessel *et al.*, 2013). Similarly, a meta-analysis of only irrigated systems in Mediterranean climates also found no tillage differences (Aguilera *et al.*, 2013). Consistent with the larger van Kessel *et al.* (2013) analysis, however, Aguilera *et al.* (2013) also noted that NT/RT increased N₂O emissions by 72% compared to CT in experiments lasting <1 year. In the present study, the 5-year mean soil N₂O emissions after 10 years of management was 22% lower in NT (2.3 ± 0.3 kg N₂O-N ha⁻¹ yr⁻¹) compared to CT (3.3 ± 0.4 kg N₂O-N ha⁻¹ yr⁻¹).

Soil N₂O emissions have been predicted to be greater in NT compared to CT due to higher bulk densities leading to wetter, less aerated soils (Linn & Doran, 1984; Baggs *et al.*, 2003; Rochette, 2008), but emissions have also been predicted to decrease in NT because of improved soil structure and cooler soil temperatures (Snyder *et al.*, 2009; Venterea *et al.*, 2011; Sainju *et al.*, 2012; Guzman *et al.*, 2015). Although CT soils were generally warmer and drier than NT in this study, we found that greater emissions resulted from more inter-annual variability in CT responses. For two of five crop-years, soil N₂O emissions nearly doubled whereas emissions during the other 3 years were similar to NT. While it is not clear why emissions were high during those 2 years, the temporal variability observed in N₂O emissions between years as well as during the year is an important factor to consider in accurately quantifying management effects on soil GHG fluxes. In addition, greater variability in weather, particularly the significant increase in mean annual air temperature and higher rainfall variability during the last 5 years of this study, approximate climate changes that have been predicted for the central Great Plains region of the United States (Bathke *et al.*, 2014). Thus, management responses here may provide some insight into potential impacts of projected climate changes on intensively managed agroecosystems.

The high rate of stover removal used in this study decreased both area-scaled and yield-scaled annual soil N₂O emissions compared to no removal, likely due to a decrease in available C and N substrate as well as altered environmental conditions. Limited data on the effect of crop residue removal on soil N₂O emissions in warm temperate irrigated corn systems suggest that residue removal effects depend on residue quality (e.g., lignin, polyphenol, C : N ratio; Liu *et al.*, 2011) and that emissions are expected to decrease when residues are removed compared to retained (Heller *et al.*, 2010; Liu *et al.*, 2011; Huang *et al.*, 2013). Other rainfed studies in warmer climates have also noted lower soil N₂O emissions when crop residue is removed (Jin *et al.*, 2014; Guzman *et al.*, 2015). In contrast, rainfed studies from colder climates have shown that stover removal can increase soil N₂O emissions (Congreaves *et al.*, 2016; Lehman & Osborne, 2016), possibly due to warmer soil temperatures or changes in moisture/aeration conditions that stimulate denitrifier activity. In a meta-analysis of 28 crop residue removal studies of mostly nonirrigated systems, Chen *et al.* (2013) found that soil N₂O emissions were positively correlated with the amount of residue returned to the soil and that residue-derived emissions were comparable to those from N fertilizer addition. Trends observed by Chen *et al.* (2013) are consistent with IPCC estimates that 1% of

both synthetic fertilizer N and crop residue N are lost to the atmosphere as N_2O (IPCC, 2007). Thus, residue removal would be expected to result in concomitant reductions in soil N_2O emissions.

Soil organic carbon changes

Losses of SOC occurred with stover removal and CT, indicating the importance of maintaining C input and reducing soil disturbance in maintaining SOC stocks in irrigated corn production systems. All soils lost C except for the no stover removal system, where SOC losses were not different than zero and were consistent with those previously measured on a fixed depth basis in Schmer *et al.* (2014a). Although the same SOC loss trends persisted and were somewhat greater in the later part of the study (2010–2014), no management effects were detected during this time. Greater SOC loss rates later in the study were unexpected, as rates of SOC change are predicted to decrease with time under long-term treatment (Stewart *et al.*, 2007). Change values for SOC, however, were highly variable during this period, perhaps due to variability in sampling and/or weather conditions. Overall, stover removal resulted in a 9% loss of SOC after 14 years of management (2001–2014). This is consistent with previous reports from this site and a similar nearby irrigated no-till continuous corn site showing measured and modeled 10-year SOC losses (2001–2010) in all management systems, with significant SOC losses in the surface 30 cm of soil (Liska *et al.*, 2014; Schmer *et al.*, 2014a; Wienhold *et al.*, 2015).

Stover removal in other irrigated corn systems has also resulted in surface soil C losses (Biau *et al.*, 2013; Kenney *et al.*, 2013; Halvorson & Stewart, 2015). Although increasing crop rotation complexity generally enhances SOC storage, limited SOC changes occur when continuous corn is converted to a corn–soybean rotation (West & Post, 2002). Thus, similar to SOC losses observed under continuous corn, residue removal in a rainfed corn–soybean system also resulted in SOC losses (Lehman & Osborne, 2016). Other corn rotational systems (e.g., corn–wheat *Triticum aestivum* L.) have shown numerical but not statistically significant SOC losses after crop residue removal (Lemke *et al.*, 2010; Huang *et al.*, 2013). In some rainfed no-till corn studies, however, surface soils gained SOC over time regardless of stover removal level, with slower accrual when stover was removed (Follett *et al.*, 2012; Jin *et al.*, 2015; Stewart *et al.*, 2015). Tillage can interact with stover management such that stover incorporation also can lead to SOC increases, as found in a meta-analysis of European agricultural systems (Lehtinen *et al.*, 2014) and in a nearby irrigated study (Adviento-Borbe *et al.*, 2007). In other tilled systems, no changes in SOC

have been observed when stover was retained and incorporated (Blanco-Canqui & Lal, 2007; Bundy *et al.*, 2011; Powlson *et al.*, 2011; Biau *et al.*, 2013). Under NT, however, stover removal can decrease SOC or slow accrual compared to no stover removal (Follett *et al.*, 2012; Sindelar *et al.*, 2014; Stewart *et al.*, 2015). While there were no significant main or interaction effects with tillage in the present study, management systems ranked from greatest SOC loss to least SOC loss in the longer term was CT-high > NT-high > CT-none > NT-none.

In a regional assessment of stover removal across the U.S. Corn Belt, Johnson *et al.* (2014) estimated that a minimum stover return rate of $5.74 \pm 2.4 \text{ Mg ha}^{-1} \text{ yr}^{-1}$ was necessary to maintain SOC levels. In the present study, SOC losses occurred even when all stover ($\sim 8.3 \pm 0.3 \text{ Mg ha}^{-1} \text{ yr}^{-1}$) was returned to soil. Our results are consistent with expectations that irrigated systems have low potential to sequester C because high biomass production is offset by SOC losses from decomposition stimulated by irrigation (Verma *et al.*, 2005; Follett *et al.*, 2013; Schmer *et al.*, 2014a). Although surface (0–30 cm) SOC changes can occur, Schmer *et al.* (2014a) found that SOC stocks in the top 150 cm of the soil profile did not change with stover removal or tillage. While maintenance and accrual of SOC in surface soils are critical to overall soil health and fertility, further research is necessary in deeper soils to better understand how management affects terrestrial SOC dynamics and C storage potential of the full soil profile (Kravchenko & Robertson, 2011; Follett *et al.*, 2013; Schmer *et al.*, 2014a).

Global warming potential

Conservation practices (e.g., NT, stover retention) each decreased GWP compared to conventional practices (e.g., CT, stover removal) when ΔSOC values were combined with soil GHG emissions and estimated fuel and irrigation usage. Here, stover removal increased system GWP because SOC losses offset decreased soil non- CO_2 GHG emissions. An irrigated corn–wheat system in China similarly found that crop residue removal increased system GWP primarily due to the negative effect on SOC stocks (Huang *et al.*, 2013). In contrast, other researchers have found that continuous corn systems using NT have neutral or negative GWP, while CT systems are net GHG sources (Grandy *et al.*, 2006; Mosier *et al.*, 2006; Follett *et al.*, 2013). Regardless of management practice, all systems evaluated here were net GHG sources, similar to a nearby irrigated continuous corn production system which used eddy covariance techniques to determine that it was also a net GHG source (Adviento-Borbe *et al.*, 2007).

Importantly, using NT and stover conservation practices together over the long term (i.e., 14 years) did not confer any additional mitigation benefit in this irrigated continuous corn system. System GWP is expected to decrease over time as both rates of SOC change and soil GHG emissions adjust to management (Six *et al.*, 2004; Stewart *et al.*, 2007; Chen *et al.*, 2013; Lehtinen *et al.*, 2014), but variability in cropping system, management equipment/timing/history, soil type, location, weather, and the depth to which Δ SOC is measured will affect the GWP outcomes of irrigated systems globally. At the regional scale, Nebraska has the highest percentage of U.S. irrigated acres (~15%; NASS, 2014); thus, this long-term irrigated study provides a valuable empirical dataset of management impacts on crop yields, soil GHG emissions, and surface SOC stocks in the western U.S. Corn Belt. Such long-term agronomic and soil outcomes can be applied by local producers to make evidence-based management decisions that promote soil health, reduce soil erosion risks, and ensure production sustainability.

Acknowledgements

This project was supported by the USDA-ARS as part of the USDA-ARS-Resilient Economic Agricultural Practices (REAP) project and the USDA-ARS-Greenhouse gas Reduction through Agricultural Carbon Enhancement Network (GRACEnet). Additional funding provided by the North Central Regional Sun-Grant Center at South Dakota State University through a grant from the US Department of Energy – Office of Biomass Programs (award DE-FC36-05GO85041). We thank many excellent students for their field and laboratory assistance over the years and express deep gratitude to our technicians Carla Ahlschwede, Chris Bauer, Aaron Bereuter, Nate Mellor, Elizabeth Pruessner, Susan Siragusa-Ortman, Steve Swanson, Susan Wagner, and David Walla. USDA is an equal opportunity provider and employer. Mention of trade names or commercial products in this publication does not imply recommendation or endorsement by the U.S. Department of Agriculture.

References

- Adler PR, Del Grosso SJ, Parton WJ (2007) Life-cycle assessment of net greenhouse-gas flux for bioenergy cropping systems. *Ecological Applications*, **17**, 675–691.
- Adviento-Borbe MAA, Haddix ML, Binder DL, Walters DT, Dobermann A (2007) Soil greenhouse gas fluxes and global warming potential in four high-yielding maize systems. *Global Change Biology*, **13**, 1972–1988.
- Aguilera E, Lassaletta L, Sanz-Cobena A, Garnier J, Vallejo A (2013) The potential of organic fertilizers and water management to reduce N₂O emissions in Mediterranean climate cropping systems. A review. *Agriculture, Ecosystems, and Environment*, **164**, 32–52.
- Alluvione F, Halvorson AD, Del Grosso SJ (2009) Nitrogen, tillage, and crop rotation effects on carbon dioxide and methane fluxes from irrigated cropping systems. *Journal of Environmental Quality*, **38**, 2023–2033.
- Baggs EM, Stevenson M, Pihlatie M, Regar A, Cook H, Cadisch G (2003) Nitrous oxide emissions following application of residues and fertiliser under zero and conventional tillage. *Plant and Soil*, **254**, 361–370.
- Bateman EJ, Baggs EM (2005) Contributions of nitrification and denitrification to N₂O emissions from soils at different water-filled pore space. *Biology and Fertility of Soils*, **41**, 379–388.
- Bathke DJ, Oglesby RJ, Rowe CM, Wilhite DA (2014) *Understanding and Assessing Climate Change: Implications for Nebraska*. University of Nebraska-Lincoln, Lincoln, NE.
- Biau A, Santiveri F, Lloveras J (2013) Stover management and nitrogen fertilization effects on corn production. *Agronomy Journal*, **105**, 1264–1270.
- Blanco-Canqui H, Lal R (2007) Soil and crop response to harvesting corn residues for biofuel production. *Geoderma*, **141**, 355–362.
- Blanco-Canqui H, Lal R (2009) Crop residue removal impacts on soil productivity and environmental quality. *Critical Reviews in Plant Science*, **28**, 139–163.
- van Bochove E, Jones HG, Bertrand N, Prévost D (2000) Winter fluxes of greenhouse gases from snow-covered agricultural soil: intra-annual and interannual variations. *Global Biogeochemical Cycles*, **14**, 113–125.
- Bronson KF, Mosier AR (1994) Suppression of methane oxidation in aerobic soil by nitrogen fertilizers, nitrification inhibitors, and urease inhibitors. *Biology and Fertility of Soils*, **17**, 263–268.
- Bundy LG, Andraski TW, Ruark MD, Peterson AD (2011) Long-term continuous corn and nitrogen fertilizer effects on productivity and soil properties. *Agronomy Journal*, **103**, 1346–1351.
- Butterbach-Bahl K, Baggs EM, Dannenmann M, Kiese R, Zechmeister-Boltenstern S (2013) Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B*, **368**, 20130122.
- California Air Resources Board (2014) Proposed re-adoption of the Low Carbon Fuel Standard, pp. 3–38. Available at: <http://www.arb.ca.gov/regact/2015/lcfs2015/lcfs15isor.pdf> (accessed 17 August 2016).
- Cavigelli MA, Parkin T (2012) Cropland management contributions to greenhouse gas flux. In: *Managing Agricultural Greenhouse Gases: Coordinated Agricultural Research through GRACEnet to Address Our Changing Climate* (eds Liebig M, Franzluebbers A, Follett RF), pp. 129–165. Academic Press an imprint of Elsevier, Waltham, MA.
- Chapuis-Lardy L, Wrage N, Metay A, Chotte J, Bernoux M (2007) Soils, a sink for N₂O? A review. *Global Change Biology*, **13**, 1–17.
- Chen H, Li X, Hu F, Shi W (2013) Soil nitrous oxide emissions following crop residue addition: a meta-analysis. *Global Change Biology*, **19**, 2956–2964.
- Congreaves KA, Brown SE, Németh DD, Dunfield KE, Wagner-Riddle C (2016) Differences in field-scale N₂O flux linked to crop residue removal under two tillage systems in cold climates. *Global Change Biology Bioenergy*, doi: 10.1111/gcbb.12354.
- Conrad R (1996) Soil microorganisms as controllers of atmospheric trace gases (H₂, CO, CH₄, OCS, N₂O and NO). *Microbiological Reviews*, **60**, 609–640.
- Cui F, Yan G, Zhou Z, Zheng X, Deng J (2012) Annual emissions of nitrous oxide and nitric oxide from a wheat-maize cropping system on a silt loam calcareous soil in the North China Plain. *Soil Biology and Biochemistry*, **48**, 10–19.
- Davidson EA (1991) Fluxes of nitrous oxide and nitric oxide from terrestrial ecosystems. In: *Microbial Production and Consumption of Greenhouse Gases* (eds Rogers JE, Whitman WB), pp. 219–235. American Society of Microbiology, Washington, DC.
- Decock C (2014) Mitigating nitrous oxide emissions from corn cropping systems in the Midwestern U.S.: potential and data gaps. *Environmental Science and Technology*, **48**, 4247–4256.
- Del Grosso SJ, White JW, Wilson G *et al.* (2013) Introducing the GRACEnet/REAP data contribution, discovery and retrieval system. *Journal of Environmental Quality*, **42**, 1274–1280.
- Drury CF, Reynolds WD, Yang XM, McLaughlin NB, Welacky TW, Calder W, Grant CA (2012) Nitrogen source, application time and tillage effects on soil nitrous oxide emissions and corn grain yields. *Soil Science Society of America Journal*, **76**, 1268–1279.
- ERS (2015) Irrigation and Water Use. Economic Research Service. Available at: <http://www.ers.usda.gov/topics/farm-practices-management/irrigation-water-use/background.aspx> (accessed 20 August 2016).
- FAO. (2011). *The State of the World's Land and Water Resources for Food and Agriculture (SOLAW) – Managing Systems at Risk*. Food and Agriculture Organization of the United Nations, Earthscan, Rome, London.
- Follett RF, Pruessner EG (2001) Intra-laboratory carbon isotope measurements on five soils. In: *Assessment Methods for Soil Carbon* (eds Lal R, Kimble JM, Follett RF, Stewart BA), pp. 185–192. CRC Press, Boca Raton, FL.
- Follett RF, Vogel KP, Varvel GE, Mitchell RB, Kimble J (2012) Soil carbon sequestration by switchgrass and no-till maize grown for bioenergy. *BioEnergy Research*, **5**, 866–875.
- Follett RF, Jantalia CP, Halvorson AD (2013) Soil carbon dynamics for irrigated corn under two tillage systems. *Soil Science Society of America Journal*, **77**, 951–963.
- Forster P, Ramaswamy V, Artaxo P *et al.* (2007) Changes in atmospheric constituents and in radiative forcing. In: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental*

- Panel on Climate Change (eds Solomon S, Qin D, Manning M *et al.*), pp. 129–234. Cambridge University Press, Cambridge, UK and New York.
- Grandy AS, Loecke TD, Parr S, Robertson GP (2006) Long-term trends in nitrous oxide emissions, soil nitrogen, and crop yields of till and no-till cropping systems. *Journal of Environmental Quality*, **35**, 1487–1495.
- Grassini P, Cassman KG (2012) High-yield maize with large net energy yield and small global warming intensity. *Proceedings of the National Academy of Sciences*, **109**, 1074–1079.
- Guzman J, Al-Kaisi M, Parkin T (2015) Greenhouse gas emissions dynamics as influenced by corn residue removal in continuous corn system. *Soil Science Society of America Journal*, **79**, 612–625.
- Halvorson AD, Stewart CE (2015) Stover removal affects no-till irrigated corn yields, soil carbon, and nitrogen. *Agronomy Journal*, **107**, 1504–1512.
- Halvorson AD, Mosier AR, Reule CA, Bausch WC (2006) Nitrogen and tillage effects on irrigated continuous corn yields. *Agronomy Journal*, **98**, 63–71.
- Halvorson AD, Del Grosso SJ, Reule CA (2008) Nitrogen, tillage, and crop rotation effects on nitrous oxide emissions from irrigated cropping systems. *Journal of Environmental Quality*, **37**, 1337–1344.
- Heller H, Bar-Tal A, Tamir G *et al.* (2010) Effects of manure and cultivation on carbon dioxide and nitrous oxide emission from a corn field under Mediterranean conditions. *Journal of Environmental Quality*, **39**, 437–448.
- Huang T, Gao B, Christie P, Ju X (2013) Net global warming potential and greenhouse gas intensity in a double-cropping cereal rotation as affected by nitrogen and straw management. *Biogeosciences*, **10**, 7897–7911.
- Hutchinson GL, Mosier AR (1981) Improved soil cover method for field measurement of nitrous oxide fluxes. *Soil Science Society of America Journal*, **43**, 311–316.
- Jin VL, Baker JM, Johnson JM-F *et al.* (2014) Soil greenhouse gas emissions in response to corn stover removal and tillage management across the US Corn Belt. *BioEnergy Research*, **7**, 517–527.
- Jin VL, Schmer MR, Wienhold BJ *et al.* (2015) Twelve years of stover removal increases soil erosion potential without impacting yield. *Soil Science Society of America Journal*, **79**, 1169–1178.
- Johnson JMF, Novak JM, Varvel GE *et al.* (2014) Crop residue mass needed to maintain soil organic carbon levels: can it be determined? *BioEnergy Research*, **7**, 481–490.
- Karlen DL, Varvel GE, Johnson JMF *et al.* (2011) Monitoring soil quality to assess the sustainability of harvesting corn stover. *Agronomy Journal*, **103**, 288–295.
- Karlen DL, Birrell SJ, Johnson JMF *et al.* (2014) Multilocation corn stover harvest effects on crop yields and nutrient removal. *BioEnergy Research*, **7**, 528–539.
- Kaspar TC, Erbach DC, Cruse RM (1990) Corn response to seed-row residue removal. *Agronomy Journal*, **54**, 1112–1117.
- Kenney I, Blanco-Canqui H, Presley DR, Rice CW, Janssen K, Olson B (2013) Soil and crop response to stover removal from rainfed and irrigated corn. *Global Change Biology Bioenergy*, **7**, 219–230.
- van Kessel C, Venterea R, Six J, Adviento-Borbe MA, Linquist B, van Groenigen KJ (2013) Climate, duration, and N placement determine N₂O emissions in reduced tillage systems: a meta-analysis. *Global Change Biology*, **19**, 33–44.
- Kravchenko AN, Robertson GP (2011) Whole-profile soil carbon stocks: the danger of assuming too much from analyses of too little. *Soil Science Society of America Journal*, **75**, 235–240.
- Lal R, Kimble JM, Follett RF, Cole CV (1998) *The Potential of U.S. Cropland to Sequester Carbon and Mitigate the Greenhouse Effect*. Ann Arbor Press, Chelsea, MI.
- Lee J, Hopmans JW, Rolston DE, Baer SG, Six J (2009) Determining soil carbon stock changes: simple bulk density corrections fail. *Agriculture, Ecosystems and Environment*, **134**, 251–256.
- Lehman RM, Osborne SL (2016) Soil greenhouse gas emissions and carbon dynamics of a no-till, corn-based cellulosic ethanol production system. *BioEnergy Research*, **9**, 1101–1108.
- Lehtinen T, Schlatter N, Baumgarten A *et al.* (2014) Effect of crop residue incorporation on soil organic carbon and greenhouse gas emissions in European agricultural soils. *Soil Use and Management*, **30**, 524–538.
- Lemke RL, VandeBygaert AJ, Campbell CA, Lafond GP, Grant B (2010) Crop residue removal and fertilizer N: effects on soil organic carbon in a long-term crop rotation experiment on a Udic Boroll. *Agriculture, Ecosystems and Environment*, **135**, 42–51.
- Li C, Frolking S, Butterbach-Bahl K (2005) Carbon sequestration in arable soils is likely to increase nitrous oxide emissions, offsetting reductions in climate radiative forcing. *Climate Change*, **72**, 321–338.
- Linn DM, Doran JW (1984) Effect of water-filled pore space on carbon dioxide and nitrous oxide production in tilled and nontilled soils. *Soil Science Society of America Journal*, **48**, 1267–1272.
- Liska AJ, Yang H, Milner M *et al.* (2014) Biofuels from crop residue can reduce soil carbon and increase CO₂ emissions. *Nature Climate Change*, **4**, 398–401.
- Littell RC, Milliken GA, Stroup WW, Wolfinger RD, Schabenberger O (2006) *SAS for Mixed Models* (2nd edn). SAS Inst., Cary, NC.
- Liu C, Wang K, Meng S *et al.* (2011) Effects of irrigation, fertilization and crop straw management on nitrous oxide and nitric oxide emissions from a wheat–maize rotation field in northern China. *Agriculture, Ecosystems and Environment*, **140**, 226–233.
- Loughin TM (2006) Improved experimental design and analysis for long-term experiments. *Crop Science*, **46**, 2492–2502.
- Maharjan B, Venterea RT, Rosen C (2014) Fertilizer and irrigation management effects on nitrous oxide emissions and nitrate leaching. *Agronomy Journal*, **106**, 703–714.
- Mahlh SS, Lemke R (2007) Tillage, crop residue and N fertilizer effects on crop yield, nutrient uptake, soil quality and nitrous oxide gas emissions in a second 4-yr rotation cycle. *Soil and Tillage Research*, **96**, 269–283.
- Mosier AR, Halvorson AD, Reule CA, Liu XJ (2006) Net global warming potential and greenhouse gas intensity in irrigated cropping systems in northeastern Colorado. *Journal of Environmental Quality*, **35**, 1584–1598.
- Mutegi JK, Munkholm LJ, Petersen BM, Hansen EM, Petersen SO (2010) Nitrous oxide emissions and controls as influenced by tillage and crop residue management strategy. *Soil Biology and Biochemistry*, **42**, 1701–1711.
- NASS (2014) Farm and Ranch Irrigation Survey (2013) Special studies Volume 3, Part 1. USDA-National Agricultural Statistics Service, 2012 Census of Agriculture, AC-12-SS-1, U.S. Government Printing Office, Washington, DC.
- NCDC (2016) Daily/Monthly Normals (1981–2010), Mead 6 S. National Climatic Data Center. Available at: <https://www.ncdc.noaa.gov/data-access/land-based-station-data/land-based-datasets/climate-normals/1981-2010-normals-data> (accessed 1 June 2016).
- Norwood CA (1999) Water use and yield of limited-irrigated and dryland corn. *Soil Science Society of America Journal*, **64**, 365–370.
- Ogle SM, Swan A, Paustian K (2012) No-till management impacts on crop productivity, carbon input and soil carbon sequestration. *Agriculture, Ecosystems and Environment*, **149**, 37–49.
- Parkin TB (2008) Effect of sampling frequency on estimates of cumulative nitrous oxide emissions. *Journal of Environmental Quality*, **37**, 1390–1395.
- Parkin TB, Venterea RT (2010) Chamber-based trace gas flux measurements. In: *Sampling Protocols* (ed. Follett RF), pp. 3–1–3–39. Available at: [http://www.ars.usda.gov/research/GRACEnet/Chamber Based Trace Gas Flux Measurements](http://www.ars.usda.gov/research/GRACEnet/Chamber%20Based%20Trace%20Gas%20Flux%20Measurements) (accessed 1 February 2011).
- Parkin TB, Venterea RT, Hargreaves SK (2012) Calculating the detection limits of chamber-based soil greenhouse gas flux measurements. *Journal of Environmental Quality*, **41**, 705–715.
- Powlson DS, Whitmore AP, Goulding KWT (2011) Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science*, **62**, 42–55.
- Reicosky DC, Lindstrom MJ, Schumacher TE, Lobb DE, Malo DD (2005) Tillage-induced CO₂ loss across an eroded landscape. *Soil and Tillage Research*, **81**, 183–194.
- Rochette P (2008) No-till only increases N₂O emissions in poorly-aerated soils. *Soil and Tillage Research*, **101**, 97–100.
- Sainju UM, Stevens WB, Caesar-TonThat T, Liebig MA (2012) Soil greenhouse gas emissions affected by irrigation, tillage, crop rotation, and nitrogen fertilization. *Journal of Environmental Quality*, **41**, 1774–1786.
- SAS Institute (2014) *The SAS System for Windows, Version 9.3*. SAS Inst., Cary, NC.
- Schaufli G, Kitzler B, Schindlbacher A, Skiba U, Sutton MA, Zechmeister-Boltenstern S (2010) Greenhouse gas emissions from European soils under different land use: effects of soil moisture and temperature. *European Journal of Soil Science*, **61**, 683–696.
- Schmer MR, Jin VL, Wienhold BJ, Varvel GE, Follett RF (2014a) Tillage and residue management effects on soil carbon and nitrogen under irrigated continuous corn. *Soil Science Society of America Journal*, **78**, 1987–1996.
- Schmer MR, Vogel KP, Varvel GE, Follett RF, Mitchell RB, Jin VL (2014b) Energy potential and greenhouse gas emissions from bioenergy cropping systems on marginally productive cropland. *PLoS One*, **9**, e89501.
- Sims AL, Schepers JS, Olson RA, Power JF (1998) Irrigated corn yield and nitrogen accumulation response in a comparison of no-till and conventional till: tillage and surface-residue variables. *Agronomy Journal*, **90**, 630–637.
- Sindelar AJ, Coulter JA, Lamb JA, Vetsch JA (2013) Agronomic responses of continuous corn to stover, tillage, and nitrogen management. *Agronomy Journal*, **105**, 1498–1506.
- Sindelar AJ, Lamb JA, Coulter JA (2014) Short-term stover, tillage, and nitrogen management affect near-surface soil organic matter. *Soil Science Society of America Journal*, **79**, 251–260.

- Sindelar AJ, Schmer MR, Jin VL, Wienhold BJ, Varvel GE (2015) Long-term corn and soybean response to crop rotation and tillage. *Agronomy Journal*, **107**, 2241–2252.
- Six J, Ogle SM, Breidt FJ, Conant RT, Mosier AR, Paustian K (2004) The potential to mitigate global warming no-tillage management is only realized when practiced in the long run. *Global Change Biology*, **10**, 155–160.
- Snyder CS, Bruulsema TW, Jensen TL, Fixen PE (2009) Review of greenhouse gas emissions from crop production systems and fertilizer management effects. *Agriculture, Ecosystems and Environment*, **133**, 247–266.
- Soil Survey Staff (2014) "Web Soil Survey," USDA-Natural Resources Conservation Service. Available at: <http://websoilsurvey.nrcs.usda.gov> (accessed 12 February 2014).
- Stewart CE, Paustian K, Conant RT, Plante AF, Six J (2007) Soil carbon saturation: concept, evidence and evaluation. *Biogeochemistry*, **86**, 19–31.
- Stewart CE, Follett RF, Pruessner EG, Varvel GE, Vogel KP, Mitchell RB (2015) Nitrogen and harvest effects on soil properties under rainfed switchgrass and no-till corn over 9 years: implications for soil quality. *Global Change Biology Bioenergy*, **7**, 288–301.
- Teepe R, Brumme R, Beese F (2000) Nitrous oxide emissions from frozen soils under agricultural, fallow and forest land. *Soil Biology and Biochemistry*, **32**, 1807–1810.
- Venterea RT (2010) Simplified method for quantifying theoretical underestimation of chamber-based trace gas fluxes. *Journal of Environmental Quality*, **39**, 126–135.
- Venterea RT, Maharjan B, Dolan MS (2011) Fertilizer source and tillage effects on yield-scaled nitrous oxide emissions in a corn cropping system. *Journal of Environmental Quality*, **40**, 1521–1531.
- Verma SB, Cobermann A, Cassman KG *et al.* (2005) Annual carbon dioxide exchange in irrigated and rainfed maize-based agroecosystems. *Agricultural and Forest Meteorology*, **131**, 77–96.
- Wagner SW, Reicosky DC, Alessi RS (1997) Regression models for calculating gas fluxes measured with a closed chamber. *Agronomy Journal*, **89**, 279–284.
- West TO, Post WM (2002) Soil organic carbon sequestration rates by tillage and crop rotation. *Soil Science Society of America Journal*, **66**, 1930–1946.
- Wienhold BJ, Schmer MR, Jin VL, Varvel GE, Gollany H (2015) CQESTR simulated changes in soil organic carbon under residue management practices in continuous corn systems. *BioEnergy Research*, **9**, 23–30.
- Wilhelm W, Johnson J, Hatfield J, Voorhees W, Linden D (2004) Crop and soil productivity response to corn residue removal: a literature review. *Agronomy Journal*, **96**, 1–17.
- Wilhelm WW, Johnson JMF, Karlen DL, Lightle DT (2007) Corn stover to sustain soil organic carbon further constrains biomass supply. *Agronomy Journal*, **99**, 1665–1667.
- Wortmann CS, Shapiro CA, Schmer MR (2016) Residue harvest effects on irrigated, no-till corn yield and nitrogen response. *Agronomy Journal*, **108**, 384–390.
- Zhang Y, Mu Y, Zhou Y, Liu J, Zhang C (2014) Nitrous oxide emissions from maize-wheat field during 4 successive years in the North China Plain. *Biogeosciences*, **11**, 1717–1726.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Effects of tillage (a, b) and stover removal (c, d) on total soil organic carbon changes (DSOC) and global warming potential (GWP) ($\text{Mg CO}_2 \text{ eq ha}^{-1}$) for 2001–2010, 2010–2014, and the full 2001–2014 time horizon. Within each time period, treatment differences are denoted by * $P \geq 0.05$, ** $P > 0.01$, and $^{\S}P = 0.0828$. In panels (a, c), daggers (†) indicate SOC changes that are different than zero.

Table S1. Annual fossil-fuel energy requirements and emissions as carbon dioxide-equivalents for agronomic management operations.